WARNING: Keep out of reach of children



# PROFESSIONAL STRENGTH

**10** kg

- · Indoor and outdoor use.
- . Hole for fixing in place.
- Octagonal blocks give rodents more edges to get their teeth into.
- Formulated to withstand moisture, damp and heat and will not break down or leach poison.
  - · Contains non-target species deterrent.

# **READ LABEL BEFORE USE**

Ready to use rodenticide bait for the control of rats and mice in and around buildings.

**INGREDIENTS:** Contains 0.05 g/kg Brodifacoum and 0.01 g/kg bittering agent Denatonium Benzoate in the form of a bait.

WARNING: Repeated or prolonged ingestion of bait may cause reduced blood clotting and increase bruising. Designed for biocidal activity

PRECAUTIONS: Keep out of reach of children. Wear gloves and avoid skin contact. Wash hands and exposed skin after handling. Avoid contamination of any water supply with product or empty container. Do not place baits in places accessible to children, pets, domestic animals, livestock, birds or where feral animals may have access to baits. Dispose of any carcasses by wrapping in paper, placing in a plastic bag and putting in the garbage.

FIRST AID: If swallowed do not induce vomiting. Seek medical advice and have product label to hand.

**Treatment of domestic animals:** Vitamin KI is an effective antidote against accidental poisoning.

Note to Doctors and Veterinarians: Brodifacoum is a long-lasting anticoagulant compound which if ingested by humans, domestic animals or pets, can reduce the clotting ability of the blood and haemorrhage may result. Vitamin KI should be administered. If administered intravenously the injection must be given slowly.

**STORAGE:** Store tightly closed in the original container in a secure, dry, cool area away from direct sunlight and foodstuffs.

**DISPOSAL:** Before disposing, wrap any untaken or unused baits in a plastic bag and ensure container is completely empty. Do not use container for any other purpose. Dispose of in the garbage.

# **BAITING STRATEGY:**

By law this product must only be used in a bait station and as specified on the label.

- Eliminate as many alternative food sources as possible.
- Human odours may deter rodents; wear gloves when handling bait.
- Place bait station under cover, along walls or rafters and in dark sheltered areas where there are signs of rodent activity.

**For mice** - Use single bait blocks in Kiwicare Bait Stations, 5 m apart, or 2 m apart for high infestations.

For rats - Use I to 3 blocks in Kiwicare Bait Stations IO mapart, or 5 m apart for high infestations.

- Inspect baits at least every 7 days.
- For effective control continue to replace eaten bait until feeding ceases. Once rodent activity has ceased ensure untaken baits are removed and safely disposed of (see Disposal).

**NOTE:** One Feed is sufficient to kill a rat or mouse but they will continue to eat bait.

• It may take up to 2 weeks to control a significant rodent problem.

As Kiwicare Corporation Limited has no control over the storage or use of this product, no liability is accepted for accidents or damage arising from the incorrect storage or use of this product.

### Distributed by:

Kiwicare Ltd. 225 Maces Road, Christchurch, NZ Phone: (+64) 3 389 0778 Fax: 64 3 389 0669 sales@kiwicare.co.nz. www.kiwicare.co.nz

## **New Zealand Agent:**

PelGar International (NZ) Limited, PO Box 11825, Ellerslie, Auckland, New Zealand. Phone: (+64) 21 670 105

#### Registered by

Pelgar International Limited, Unit 13, Newman Lane, Alton,

Hants, GU34 2QR, UK.

Phone: (+44) 1420 80744 www.pelgar.co.uk

#### EPA Approval Number: HSR001594

Approved Maintenance Compound (Non-Dairy)- Type C. Registered pursuant to the ACVM Act 1997, No. V960I See www.foodsafety.govt.nz for registration conditions.



